arXiv:1207.0638v1 [math.CO] 3 Jul 2012

Isoperimetric Inequalities in Simplicial Complexes

Ori Parzanchevski, Ron Rosenthal and Ran J. Tessler

July 4, 2012

Abstract

In graph theory there are intimate connections between the expansion properties of a graph and the spectrum of its Laplacian. In this paper we define a notion of combinatorial expansion for simplicial complexes of general dimension, and prove that similar connections exist between the combinatorial expansion of a complex, and the spectrum of the high dimensional Laplacian defined by Eckmann. In particular, we present generalizations of the Cheeger inequalities, and the Expander Mixing lemma. As a corollary, using the work of Pach, we obtain a connection between spectral properties of complexes and Gromov's notion of geometric overlap. Using the work of Gunder and Wagner, we give an estimate for the combinatorial expansion and geometric overlap of random complexes.

1 Introduction

It is a cornerstone of graph theory that the expansion properties of a graph are intimately linked to the spectrum of its Laplacian. In particular, the discrete Cheeger inequalities [Tan84, Dod84, AM85, Alo86] relate the spectral gap of a graph to its Cheeger constant, and the Expander Mixing Lemma [AC88] relates the extremal values of the spectrum to the discrepancy in the graph (see (1.4)) and to its mixing properties.

In this paper we define a notion of expansion for simplicial complexes, which generalizes the Cheeger constant and the discrepancy in graphs. We then study its relations to the spectrum of the high dimensional Laplacian defined by Eckmann [Eck44], and present high dimensional analogues of the discrete Cheeger inequalities and the Expander Mixing Lemma.

This study is closely related to the notion of *high dimensional expanders*. A family of graphs $\{G_i\}$ with uniformly bounded degrees is said to be a family of *expanders* if their Cheeger constants $h(G_i)$ are uniformly bounded away from zero. By the discrete Cheeger inequalities (1.3), this is equivalent to having their spectral gaps $\lambda(G_i)$ uniformly bounded away from zero. Thus, combinatorial expanders and spectral expanders are equivalent notions. We refer to [HLW06, Lub12] for the general background on expanders and their applications.

It is desirable to have a similar situation in higher dimensions, but at least as of now, it is not clear what is the "right" notion of "high dimensional expander". One generalization of the Cheeger constant to higher dimensions is the notion of *cohomological expansion*, originating in [LM06, Gro10, NR12], and also in [GW12] under the name "combinatorial expansion". While in dimension one it coincides with the Cheeger constant, it was shown by Gunder and Wagner [GW12] that in general spectral expansion does not implies cohomological expansion. They show the existence, in any dimension greater than one, of complexes with complete skeletons whose spectral gaps[1] are bounded away from zero, and which have arbitrarily small cohomological expansion.

Another notion of expansion is Gromov's *geometric overlap property*, originating in [Gro10] and studied in [FGL$^+$11, MW11]. This notion was shown in [Gro10, MW11] to be related to cohomological expansion. However, even in dimension one it is not equivalent to that of expander graphs.

Our definition of expansion suggests a natural notion of "combinatorial expanders", and we show that spectral expanders with complete skeletons are combinatorial expanders, and that good enough combinatorial expanders are spectral expanders. A theorem of Pach [Pac98] shows that this notion of combinatorial expansion is also connected to the geometric overlap property.

Combinatorial Expansion and the Spectral Gap

The Cheeger constant of a finite graph $G = (V, E)$ is usually taken to be

$$\varphi(G) = \min_{\substack{A \subseteq V \\ 0<|A|<\frac{|V|}{2}}} \frac{|E(A, V \backslash A)|}{|A|}$$

where $E(A, B)$ is the set of edges with one vertex in A and the other in B. In this paper, however, we will work with the following version:

$$h(G) = \min_{0<|A|<|V|} \frac{|V|\,|E(A, V\backslash A)|}{|A|\,|V\backslash A|}. \tag{1.1}$$

Since $\varphi(G) \le h(G) \le 2\varphi(G)$, defining expanders by φ or by h is equivalent.

The *spectral gap* of G, denoted $\lambda(G)$, is the second smallest eigenvalue of the *Laplacian* $\Delta^+ : \mathbb{R}^V \to \mathbb{R}^V$, which is defined by

$$(\Delta^+ f)(v) = \deg(v) f(v) - \sum_{w \sim v} f(w). \tag{1.2}$$

The discrete Cheeger inequalities [Tan84, Dod84, AM85, Alo86] relate the Cheeger constant and the spectral gap, by

$$\frac{h^2(G)}{8k} \le \lambda(G) \le h(G), \tag{1.3}$$

where k is the maximal degree of a vertex in G.[2]

Let X be an (abstract) simplicial complex with vertex set V. This means that X is a collection of subsets of V, called *cells* (and also *simplexes*, *faces*, or *hyperedges*), which is closed under taking subsets, i.e., if $\sigma \in X$ and $\tau \subseteq \sigma$, then $\tau \in X$. The *dimension* of a cell σ is $\dim\sigma = |\sigma| - 1$, and X^j denotes the set of cells of dimension j. The dimension of X is the maximal dimension of a cell in it. The *degree* of a j-cell (a cell of dimension j) is the number of $(j+1)$-cells in which it is contained. We shall occasionally add the assumption that X has a complete skeleton, by which we mean that $X^j = \binom{V}{j+1}$ for $j < \dim X$ (here $\binom{V}{j} = \{S \subseteq V \mid |S| = j\}$).

We define the following generalization of the Cheeger constant:

---

[1] The spectral gap of a complex is defined in Section 2.1.

[2] For φ they are given by $\frac{\varphi^2(G)}{2k} \le \lambda(G) \le 2\varphi(G)$.

Definition 1.1. For a finite d-dimensional complex X with vertex set V,

$$h(X) = \min_{V=\coprod_{i=0}^{d} A_i} \frac{|V| \cdot |F(A_0, A_1, \ldots, A_d)|}{|A_0| \cdot |A_1| \cdot \ldots \cdot |A_d|},$$

where the minimum is over all partitions of V into nonempty sets $A_0, \ldots, A_d$, and $F(A_0, \ldots, A_d)$ denotes the set of d-dimensional cells with one vertex in each A_i.

For $d = 1$, this coincides with the Cheeger constant of a graph (1.1). To formulate an analogue of the Cheeger inequalities, we need a general dimension analogue of the spectral gap. Such an analogue is provided by the work of Eckmann on discrete Hodge theory [Eck44]. In order to give the definition we shall need more terminology, and we defer this to Section 2.1. The basic idea, however, is the same as for graphs, namely, it is the smallest nontrivial eigenvalue of a suitable Laplacian operator. The following theorem then generalizes the Cheeger inequalities to higher dimensions:

Theorem 1.2 (Cheeger Inequalities)**.** *Let X be a finite d-dimensional simplicial complex with spectral gap $\lambda(X)$.*

(1) If X is has a complete skeleton, then

$$\lambda(X) \le h(X).$$

(2) Let k be the maximal degree of a $(d-1)$-cell in X. Then

$$\frac{d\left(1 - \frac{d-1}{|V|}\right)^2}{8k} h^2(X) - (d-1)k \le \lambda(X)$$

(here the skeleton is not necessarily complete).

The proof appears in Section 4.1.

Remarks.

(1) Since graphs have by definition a complete skeleton, this coincides with the Cheeger inequalities for graphs (1.3).

(2) For $d > 1$ this only gives a lower affine bound on the spectral gap. Even if this bound can be improved, the affine part cannot be completely removed, not even for complexes with a complete skeleton: a counterexample is provided by the minimal triangulation of the Möbius strip (Figure 1).



Figure 1: A triangulation of the Möbius strip which has a complete skeleton, and for which $h = 1\frac{1}{4}$, $\lambda = 0$.

In [LM06] Linial and Meshulam introduced the following model for random simplicial complexes: for a given $p = p(n) \in (0,1)$, $X(d,n,p)$ is a d-dimensional simplicial complex on n vertices, with a complete skeleton, and with every d-cell being included independently with probability p. Relying on the analysis of the spectrum of $X(d,n,p)$ in [GW12], we show the following:

Corollary 1.3. *Fix $c > 0$, and observe $X = X\left(d,n,\frac{C\cdot\log n}{n}\right)$. For every $C \geq C_0(c,d)$ there exists a constant $H = H(C) > 0$ satisfying $\lim\limits_{C\to\infty}\frac{H}{C} = 1$ such that for $n \geq n_0(c,d)$*

$$\operatorname{Prob}\left(h(X) \geq H\cdot\log n\right) \geq 1 - n^{-c}.$$

The proof appears in Section 4.4, as part of Corollary 4.2.

Mixing, Discrepancy and Geometric Overlap

The Cheeger inequalities describe the expansion along the various partitions of a graph, in terms of its spectral gap. However, the spectral gap alone does not suffice to determine the expansion between small sets of vertices, and the mixing quality of the graph. For example, the bipartite Ramanujan graphs constructed in [LPS88] are regular graphs with very large spectral gaps, which are bipartite. This means that there they contain disjoint sets $A, B \subseteq V$ of size $\frac{|V|}{4}$ with $E(A,B) = 0$, and also that the simple random walk on them does not converge.

An understanding of small expansion and of mixing can be obtained by observing not only the smallest nontrivial eigenvalue of the Laplacian, but also the largest one[3]. In particular, the so-called Expander Mixing Lemma (see e.g., [AC88]) states that for a k-regular graph $G = (V,E)$, and $A, B \subseteq V$,

$$\left| |E(A,B)| - \frac{k|A||B|}{|V|} \right| \leq \rho\cdot\sqrt{|A||B|}, \tag{1.4}$$

where ρ is the maximal absolute value of a nontrivial eigenvalue of $kI - \Delta^+$.

The deviation of $|E(A,B)|$ from its expected value $p|A||B|$, where $p = \frac{k}{|V|} \approx |E|/\binom{|V|}{2}$ is the edge density, is called the *discrepancy* of A and B. This is a measure of quasi-randomness in a graph, a notion closely related to expansion (see e.g. [Chu97]). In a similar fashion, we call the deviation

$$\left| |F(A_0,\ldots,A_d)| - \frac{|X^d|}{\binom{|V|}{d+1}}\cdot|A_0|\cdot\ldots\cdot|A_d| \right|$$

the discrepancy of $A_0,\ldots,A_d$, and the following theorem bounds the discrepancy in complexes in an analogous way to the Expander Mixing Lemma:

Theorem 1.4 (Mixing Lemma)**.** *Let $\alpha \in \mathbb{R}$. If X is a d-dimensional complex with a complete skeleton, then for any disjoint sets of vertices $A_0,\ldots,A_d$ one has*

$$\left| |F(A_0,\ldots,A_d)| - \frac{\alpha\cdot|A_0|\cdot\ldots\cdot|A_d|}{|V|} \right| \leq \rho_\alpha\cdot\left(|A_0|\cdot\ldots\cdot|A_d|\right)^{\frac{d}{d+1}},$$

where ρ_α is the maximal absolute value of a nontrivial eigenvalue of $\alpha I - \Delta^+$.

Here Δ^+ is the Laplacian of X, which is defined in Section 2. The proof, and a formal definition of ρ_α, appear in Section 4.2.

---

[3] Graphs having both of them bounded are referred to as "two-sided expanders" in [Tao11].

Remark. For k-regular graphs, taking $\alpha = k$ gives the Expander Mixing Lemma (1.4).

As mentioned above, the mixing quality of a graph is also reflected by the behavior of the random walk on it. In a future paper [PR12] we suggest a generalization of the notion of random walk to high dimensional simplicial complexes, and study its connection to the spectral properties of the complex.

Let $G = (V, E)$ be a graph with a high Cheeger constant. Given a mapping $\varphi : V \to \mathbb{R}$, there exists a point $x \in \mathbb{R}$ which is covered by many edges in the linear extension of φ to E. This observation led Gromov to define the *geometric overlap* of a complex [Gro10]:

Definition 1.5. Let X be a d-dimensional simplicial complex. The overlap of X is defined by

$$\operatorname{overlap}(X) = \min_{\varphi: V \to \mathbb{R}^d} \max_{x \in \mathbb{R}^d} \frac{\#\left\{\sigma \in X^d \,\middle|\, x \in \operatorname{conv}\left\{\varphi(v) \,|\, v \in \sigma\right\}\right\}}{\left|X^d\right|}.$$

In other words, X has overlap $\geq \varepsilon$ if for every simplicial mapping of X into $\mathbb{R}^d$ (a mapping induced linearly by the images of the vertices), there is a point in $\mathbb{R}^d$ covered by at least an ε-fraction of the d-cells of X.

A theorem of Pach [Pac98], together with Theorem 1.4 yield a relation between the spectrum of the Laplacian and the overlap property. Here $\lambda(X) = \lambda_{\min}(X)$ and $\lambda_{\max}(X)$ denote the minimal and maximal nontrivial eigenvalues of the Laplacian on X.

Corollary 1.6. *There exists $\varepsilon_d > 0$ with the following property: for every $0 < \varepsilon < \varepsilon_d$ there exists $\delta > 0$, such that if X is a d-dimensional complex with a complete skeleton satisfying*

$$\frac{\lambda_{\min}(X)}{\lambda_{\max}(X)} > 1 - \varepsilon,$$

then

$$\operatorname{overlap}(X) > \delta \cdot \frac{|V|^d}{\left|X^d\right|} \cdot \lambda_{\max}(X).$$

The proof appears in Section 4.3. Returning to the $X(d, n, p)$ model, we obtain the following:

Corollary 1.7. *Fix $c > 0$. There exists a constant $\mathcal{O} = \mathcal{O}_d > 0$, such that for $C \geq C_0(c, d)$ and $n \geq n_0(c, d)$*

$$\operatorname{Prob}\left(\operatorname{overlap}\left(X\left(d, n, \frac{C \cdot \log n}{n}\right)\right) > \mathcal{O}\right) \geq 1 - n^{-c}.$$

Again, this is a part of Corollary 4.2, which is proved in Section 4.4.

The structure of the paper is as follows: in section 2 we present the basic definitions relating to simplicial complexes and their spectral theory. Section 3 is devoted to proving basic properties regarding the high dimensional Laplacians. In section 4 we prove the theorems and corollaries stated in the introduction, and section 5 lists some open questions.

Acknowledgements

The authors would like to thank Alex Lubotzky for initiating our study of expansion via spectral properties of complexes. We would also like to express our gratitude for the suggestions made to us by Noam Berger, Konstantin Golubev, Gil Kalai, Nati Linial, Doron Puder, Doron Shafrir, Uli Wagner, and Andrej Zuk. We are thankful for the support of the ERC.

2 Notations and definitions

Throughout this paper X denotes a finite d-dimensional simplicial complex with vertex set V. For $-1 \le j \le d$ we denote by X^j the set of j-cells of X. In particular, we have $X^{-1} = \{\varnothing\}$. For $j \ge 1$, every j-cell $\sigma = \left\{\sigma_0, \dots, \sigma_j\right\}$ has two possible orientations, corresponding to the possible orderings of its vertices, up to an even permutation (1-cells and the empty cell have only one orientation). We denote an oriented cell by square brackets, and a flip of orientation by an overbar. For example, one orientation of $\sigma = \{x, y, z\}$ is $[x, y, z]$, which is the same as $[y, z, x]$ and $[z, x, y]$. The other orientation of σ is $\overline{[x, y, z]} = [y, x, z] = [x, z, y] = [z, y, x]$. We denote by $X^j_\pm$ the set of oriented j-cells (so that $\left|X^j_\pm\right| = 2\left|X^j\right|$ for $j \ge 1$ and $X^j_\pm = X^j$ for $j = -1, 0$).

We now describe the *discrete Hodge theory* due to Eckmann [Eck44]. This is a discrete analogue of Hodge theory in Riemannian geometry, but in contrast, the proofs of the statements are all exercises in finite-dimensional linear algebra.

The space of *j-forms on X*, denoted by $\Omega^j(X)$, is the vector space of skew-symmetric functions on oriented j-cells:

$$\Omega^j = \Omega^j(X) = \left\{ f : X^j_\pm \to \mathbb{R} \,\middle|\, f(\overline{\sigma}) = -f(\sigma) \ \ \forall \sigma \in X^j_\pm \right\}.$$

In particular, Ω^0 is the space of functions on V, and $\Omega^{-1} = \mathbb{R}^{\{\varnothing\}}$ can be identified in a natural way with $\mathbb{R}$. We endow each Ω^i with the inner product

$$\langle f, g\rangle = \sum_{\sigma \in X^i} f(\sigma) g(\sigma) \tag{2.1}$$

(note that $f(\sigma) g(\sigma)$ is well defined even without choosing an orientation for σ).

For a cell σ (either oriented or non-oriented) and a vertex v, we denote $v \sim \sigma$ if $v \notin \sigma$ and $\{v\} \cup \sigma$ is a cell in X (here we ignore the orientation of σ). If $\sigma = \left[\sigma_0, \dots, \sigma_j\right]$ is oriented and $v \sim \sigma$, then $v\sigma$ denotes the oriented $(j+1)$-cell $\left[v, \sigma_0, \dots, \sigma_j\right]$.

The j^{th} *boundary operator* $\partial_j : \Omega^j \to \Omega^{j-1}$ is

$$\left(\partial_j f\right)(\sigma) = \sum_{v \sim \sigma} f(v\sigma).$$

The sequence $\left(\Omega^j, \partial_j\right)$ is a chain complex, i.e., $\partial_{j-1}\partial_j = 0$ for all j, and one denotes

$$\begin{aligned} Z_j &= \ker \partial_j && j\text{-cycles} \\ B_j &= \operatorname{im} \partial_{j+1} && j\text{-boundaries} \\ H_j &= {}^{Z_j}\!/_{B_j} && \text{the } j^{\text{th}} \text{ homology of } X \ (\text{over } \mathbb{R}). \end{aligned}$$

The adjoint of ∂_j w.r.t. the inner product (2.1) is the *co-boundary operator* $\partial^*_j : \Omega^{j-1} \to \Omega^j$ given by

$$\left(\partial^*_j f\right)(\sigma) = \sum_{i=0}^{j} (-1)^i f(\sigma \backslash \sigma_i)$$

where $\sigma \backslash \sigma_i = \left[\sigma_0, \sigma_1, \dots, \sigma_{i-1}, \sigma_{i+1}, \dots \sigma_j\right]$. Here the standard terms are

$$\begin{aligned} Z^j &= \ker \partial^*_{j+1} = B_j^\perp && \text{closed } j\text{-forms} \\ B^j &= \operatorname{im} \partial^*_j = Z_j^\perp && \text{exact } j\text{-forms} \\ H^j &= {}^{Z^j}\!/_{B^j} && \text{the } j^{\text{th}} \text{ cohomology of } X \ (\text{over } \mathbb{R}). \end{aligned}$$

The *upper, lower, and full Laplacians* $\Delta^+, \Delta^-, \Delta : \Omega^{d-1} \to \Omega^{d-1}$ are defined by

$$\Delta^+ = \partial_d \partial_d^*, \qquad \Delta^- = \partial_{d-1}^* \partial_{d-1}, \qquad \Delta = \Delta^+ + \Delta^-,$$

respectively[4]. All the Laplacians decompose (as a direct sum of linear operators), with respect to the orthogonal decompositions $\Omega^{d-1} = B^{d-1} \oplus Z_{d-1} = B_{d-1} \oplus Z^{d-1}$. In addition, $\ker \Delta^+ = Z^{d-1}$ and $\ker \Delta^- = Z_{d-1}$.

The space of *harmonic* $(d-1)$*-forms* on X is $\mathcal{H}_{d-1} = \ker \Delta$. If $f \in \mathcal{H}_{d-1}$ then

$$0 = \langle \Delta f, f \rangle = \langle \partial_{d-1} f, \partial_{d-1} f \rangle + \langle \partial_d^* f, \partial_d^* f \rangle$$

which shows that $\mathcal{H}_{d-1} = Z^{d-1} \cap Z_{d-1}$. This gives the so-called *discrete Hodge decomposition*

$$\Omega^{d-1} = B^{d-1} \oplus \mathcal{H}_{d-1} \oplus B_{d-1}.$$

In particular, this shows that the space of harmonic forms can be identified with the cohomology of X:

$$H^{d-1} = \frac{Z^{d-1}}{B^{d-1}} = \frac{B_{d-1}^{\perp}}{B^{d-1}} = \frac{B^{d-1} \oplus \mathcal{H}_{d-1}}{B^{d-1}} \cong \mathcal{H}_{d-1}.$$

The same holds for the homology of X, giving

$$H^{d-1} \cong \mathcal{H}_{d-1} \cong H_{d-1}. \tag{2.2}$$

For comparison, the original Hodge decomposition states that for a Riemannian manifold M and $0 \le j \le \dim M$, there is an orthogonal decomposition

$$\Omega^j(M) = d\left(\Omega^{j-1}(M)\right) \oplus \mathcal{H}^j(M) \oplus \delta\left(\Omega^{j+1}(M)\right)$$

where Ω^j are the smooth j-forms on M, d is the exterior derivative, δ its Hodge dual, and $\mathcal{H}^j$ the smooth harmonic j-forms on M. An in the discrete case, this gives an isomorphism between the j^{th} de-Rham cohomology of M and the space of harmonic j-forms on it.

Example. For $j = 0$, Z^0 consists of the locally constant functions (functions constant on connected components); B^0 consists of the constant functions; Z_0 of the functions whose sum vanishes, and B_0 of the functions whose sum on each connected component vanishes.

For $j = 1$, Z^1 are the forms whose sum along the boundary of every triangle in the complex vanishes; in B^1 lie the forms whose sum along every closed path vanishes; Z_1 are the *Kirchhoff forms*, also known as *flows*, those for which the sum over all edges incident to a vertex, oriented inward, is zero; and B_1 are the forms spanned (over $\mathbb{R}$) by oriented boundaries of triangles in the complex.

The chain of simplicial forms in dimensions -1 to 2 is depicted in Figure 2.

---

[4] More generally, one can define the j^{th} *lower Laplacian* $\Delta_j^- : \Omega^j \to \Omega^j$ by $\Delta_j^- = \partial_j^* \partial_j$, and similarly for Δ_j^+ and Δ_j. For our purposes, Δ_{d-1}^-, Δ_{d-1}^+ and Δ_{d-1} are the relevant ones.

$$\begin{tikzcd}
 & H^0 \arrow[r, phantom, "\cong"] & \mathcal{H}_0 \arrow[r, phantom, "\cong"] & H_0 & H^1 \arrow[r, phantom, "\cong"] & \mathcal{H}_1 \arrow[r, phantom, "\cong"] & H_1 & H^2 & \\
 & \underset{\substack{\text{locally}\\ \text{constant}}}{Z^0} \arrow[u, two heads] \arrow[r] \arrow[dr, hook] & \cap \arrow[u, dash] & \underset{\substack{\text{sum}\\ \text{zero}}}{Z_0} \arrow[u, two heads] \arrow[l] \arrow[dl, hook] & \underset{\substack{\text{sum zero along}\\ \text{triangle boundaries}}}{Z^1} \arrow[u, two heads] \arrow[r] \arrow[dr, hook] & \cap \arrow[u, dash] & \underset{\text{Kirchhoff}}{Z_1} \arrow[u, two heads] \arrow[l] \arrow[dl, hook] & Z^2 \arrow[u, two heads] \arrow[dr, hook] & \\
\mathbb{R} \cong \Omega^{-1} \arrow[r, shift right, "\partial_0^*"'] \arrow[d, dash, "="] & & \Omega^0 \arrow[ll, shift right, "\partial_0"'] \arrow[rrr, shift right, "\partial_1^*"'] \arrow[dll] \arrow[dl] \arrow[dr] \arrow[drr] & & & \Omega^1 \arrow[lll, shift right, "\partial_1"'] \arrow[rrr, shift right, "\partial_2^*"'] \arrow[dll] \arrow[dl] \arrow[dr] \arrow[drr] & & & \Omega^2 \arrow[lll, shift right, "\partial_2"'] \arrow[dll] \arrow[dl] \\
B_{-1} \arrow[r, dash, "\cong"] & \underset{\text{constant}}{B^0} \arrow[uu, hook] & & \underset{\substack{\text{sum zero}\\ \text{on components}}}{B_0} \arrow[uu, hook] \arrow[r, dash, "\cong"] & \underset{\substack{\text{sum zero}\\ \text{along cycles}}}{B^1} \arrow[uu, hook] & & \underset{\substack{\text{span of}\\ \text{triangle boundaries}}}{B_1} \arrow[uu, hook] \arrow[r, dash, "\cong"] & B^2 \arrow[uu, hook] &
\end{tikzcd}$$

Figure 2: The lowermost part of the chain complex of simplicial forms.

2.1 Definition of the Spectral Gap

Every graph has a "trivial zero" in the spectrum of its upper Laplacian, corresponding to the constant functions. There can be more zeros in the spectrum, and these encode information about the graph (its connectedness), while the first one does not. Similarly, for a d-dimensional complex, the space B^{d-1} is always in the kernel of the upper Laplacian, and considered to be its "trivial zeros". The existence of more zeros indicates a nontrivial $(d-1)$-cohomology, since it means that $B^{d-1} \subsetneq \ker\Delta^+ = Z^{d-1}$. Since $\left(B^{d-1}\right)^{\perp} = Z_{d-1}$, this leads to the following definition:

Definition 2.1. The spectral gap of a d-dimensional complex X, denoted $\lambda(X)$, is the minimal eigenvalue of the upper or the full Laplacian on $(d-1)$-cycles:

$$\lambda(X) = \min \operatorname{Spec}\left(\Delta\Big|_{Z_{d-1}}\right) = \min \operatorname{Spec}\left(\Delta^+\Big|_{Z_{d-1}}\right)$$

(the equality is since $\Delta\Big|_{Z_{d-1}} \equiv \Delta^+\Big|_{Z_{d-1}}$.)

The following proposition gives two more characterizations of the spectral gap, and more explicit ones for complexes with a complete skeleton appear in Proposition 3.3.

Proposition 2.2. *Let X be a d-dimensional complex, with*

$$\operatorname{Spec}\Delta^+ = \left\{\lambda_0 \leq \lambda_1 \leq \ldots \leq \lambda_{|X^{d-1}|-1}\right\}.$$

(1) If $\beta_j = \dim H_j$ is the j^{th} (reduced) Betti number of X, then

$$\lambda(X) = \lambda_r \qquad \text{where} \qquad r = \left(\left|X^{d-1}\right| - \beta_{d-1}\right) - \left(\left|X^d\right| - \beta_d\right).$$

(2) $\lambda(X)$ is the minimal nonzero eigenvalue of Δ^+, unless X has a nontrivial $(d-1)^{\text{th}}$-homology, in which case $\lambda(X) = 0$.

Remark. For a graph $G = (V, E)$, Definition 2.1 states that $\lambda(G)$ is the minimal eigenvalue of the Laplacian on a function which sums to zero. By Proposition 2.2 (1) we have $\lambda(G) = \lambda_r$, where $r = |V| - |E| - \beta_0 + \beta_1$. Since $\beta_0 + 1$ is the number of connected components in G, and β_1 is the number of cycles in G, by Euler's formula

$$r = |V| - |E| - \beta_0 + \beta_1 = \chi(G) - (\chi(G) - 1) = 1$$

and therefore $\lambda(G) = \lambda_1$. From (2) in Proposition 2.2 we obtain that $\lambda(G)$ is the minimal nonzero eigenvalue of G's Laplacian if G is connected, and zero otherwise.

Proof. Since Δ^+ decomposes w.r.t. $\Omega^{d-1} = B^{d-1} \oplus Z_{d-1}$, and $\Delta^+\big|_{B^{d-1}} \equiv 0$, the spectrum of Δ^+ consists of $r = \dim B^{d-1}$ zeros, followed by the spectral gap. By (2.2),

$$H_{d-1} \cong \mathcal{H}_{d-1} = Z^{d-1} \cap Z_{d-1} = \ker \Delta^+\Big|_{Z_{d-1}}$$

so that $\lambda(X) = 0$ if and only if $H_{d-1} \neq 0$, i.e. X has a nontrivial $(d-1)^{\text{th}}$-homology. This also shows that if $H_{d-1} = 0$, then $\lambda(X)$ is the smallest nonzero eigenvalue of Δ^+. Finally, to compute $r = \dim B^{d-1}$, we observe that

$$\begin{aligned}\dim B^{j-1} &= \dim Z^{j-1} - \dim H^{j-1} = \operatorname{null} \partial_j^* - \beta_{j-1} \\ &= \dim \Omega^{j-1} - \operatorname{rank} \partial_j^* - \beta_{j-1} = \left|X^{j-1}\right| - \dim B^j - \beta_{j-1}\end{aligned}$$

and therefore

$$\begin{aligned}r = \dim B^{d-1} &= \left|X^{d-1}\right| - B^d - \beta_{d-1} = \left|X^{d-1}\right| - \left(\left|X^d\right| - \dim B^{d+1} - \beta_d\right) - \beta_{d-1} \\ &= \left(\left|X^{d-1}\right| - \beta_{d-1}\right) - \left(\left|X^d\right| - \beta_d\right).\end{aligned}$$

□

3 Properties of the Laplacians

In this section we begin the study of the Laplacians and their spectra, obtaining results which will be used for the proofs of the main theorems. We begin by writing the Laplacians in a more explicit form.

For the upper Laplacian, if $f \in \Omega^{d-1}$ and $\sigma \in X^{d-1}$, then

$$\begin{aligned}(\Delta^+ f)(\sigma) &= \sum_{v \sim \sigma} (\partial_{d-1}^* f)(v\sigma) = \sum_{v \sim \sigma} \sum_{i=0}^{d} (-1)^i f(v\sigma \backslash (v\sigma)_i) \\ &= \sum_{v \sim \sigma} f(\sigma) - \sum_{i=0}^{d-1} (-1)^i f(v\sigma \backslash \sigma_i) \\ &= \deg(\sigma) f(\sigma) - \sum_{v \sim \sigma} \sum_{i=0}^{d-1} (-1)^i f(v\sigma \backslash \sigma_i), \qquad (3.1)\end{aligned}$$

where we recall that $\deg(\sigma)$ is the number of d-cells containing σ. For $\sigma, \sigma' \in X_{\pm}^{d-1}$ we denote $\sigma' \sim \sigma$ if there exists an oriented d-cell τ such that both σ and $\overline{\sigma'}$ are faces of τ with the orientation induced from it. Namely, if $\sigma = [\sigma_0, , \ldots, \sigma_{d-1}]$, $\sigma' = [\sigma_0, \ldots, \sigma_{i-1}, \sigma_i', \sigma_{i+1}, \ldots, \sigma_{d-1}]$, and $\{\sigma_0, \ldots, \sigma_{i-1}, \sigma_i, \sigma_i', \sigma_{i+1}, \ldots, \sigma_{d-1}\} \in X^d$. Using this notation we can express Δ^+ more elegantly as

$$(\Delta^+ f)(\sigma) = \deg(\sigma) f(\sigma) - \sum_{\sigma' \sim \sigma} f(\sigma'). \qquad (3.2)$$

For the lower Laplacian we have

$$\left(\Delta^{-}f\right)(\sigma)=\sum_{i=0}^{d-1}(-1)^{i}\left(\partial_{d-1}f\right)(\sigma\backslash\sigma_{i})=\sum_{i=0}^{d-1}(-1)^{i}\sum_{v\sim\sigma\backslash\sigma_{i}}f\left(v\sigma\backslash\sigma_{i}\right). \tag{3.3}$$

The following straightforward claim bounds the spectrum of the upper Laplacian:

Claim 3.1. *The spectrum of Δ^{+} lies in the interval $\left[0,(d+1)k\right]$, where k is the maximal degree in X.* □

3.1 Complexes with a complete skeleton

Complexes with a complete skeleton appear to be particularly well behaved, in comparison with the general case. The following proposition lists some observations regarding their Laplacians. These will be used in the proofs of the main theorems, and also to obtain simpler characterizations of the spectral gap in this case.

Proposition 3.2. *Let X be a d-dimensional complex with a complete skeleton.*

(1) If $\overline{X}$ is the complement complex of X, i.e., $\overline{X}^{d-1}=X^{d-1}=\binom{V}{d}$ and $\overline{X}^{d}=\binom{V}{d+1}\backslash X^{d}$, then

$$\Delta_{\overline{X}}^{+}=|V|\cdot I-\Delta_{X}. \tag{3.4}$$

(2) The spectrum of Δ lies in the interval $\left[0,|V|\right]$.

(3) The lower Laplacian satisfies

$$\Delta^{-}=|V|\cdot\mathbb{P}_{B^{d-1}} \tag{3.5}$$

where $\mathbb{P}_{B^{d-1}}$ is the orthogonal projection onto B^{d-1}.

Proof. By the completeness of the skeleton, the lower Laplacian (see (3.3)) can be written as

$$\begin{aligned}\left(\Delta^{-}f\right)(\sigma)&=\sum_{i=0}^{d-1}(-1)^{i}\sum_{v\sim\sigma\backslash\sigma_{i}}f\left(v\sigma\backslash\sigma_{i}\right)=\sum_{i=0}^{d-1}(-1)^{i}\sum_{v\notin\sigma\backslash\sigma_{i}}f\left(v\sigma\backslash\sigma_{i}\right)\\&=d\cdot f(\sigma)+\sum_{v\notin\sigma}\sum_{i=0}^{d-1}(-1)^{i}f\left(v\sigma\backslash\sigma_{i}\right).\end{aligned}$$

To show (1) we observe that $v\sim\sigma$ in $\overline{X}$ iff $v\notin\sigma$ and $v\nsim\sigma$ (in X), so that

$$\begin{aligned}\left(\Delta_{X}f+\Delta_{\overline{X}}^{+}f\right)(\sigma)&=\left(\Delta_{X}^{-}f\right)(\sigma)+\left(\Delta_{X}^{+}f\right)(\sigma)+\left(\Delta_{\overline{X}}^{+}f\right)(\sigma)\\&=d\cdot f(\sigma)+\sum_{v\notin\sigma}\sum_{i=0}^{d-1}(-1)^{i}f\left(v\sigma\backslash\sigma_{i}\right)\\&\quad+\deg(\sigma)f(\sigma)-\sum_{v\sim\sigma}\sum_{i=0}^{d-1}(-1)^{i}f\left(v\sigma\backslash\sigma_{i}\right)\\&\quad+\left(|V|-d-\deg(\sigma)\right)f(\sigma)-\sum_{\substack{v\notin\sigma\\v\nsim\sigma}}\sum_{i=0}^{d-1}(-1)^{i}f\left(v\sigma\backslash\sigma_{i}\right)=|V|f(\sigma).\end{aligned}$$

From (1) we conclude that $\operatorname{Spec}\Delta_{\overline{X}}^{+} = \left\{ |V| - \gamma \,\middle|\, \gamma \in \operatorname{Spec}\Delta_X \right\}$, and since Δ_X and $\Delta_{\overline{X}}^{+}$ are positive semidefinite, (2) follows. To establish (3), recall that $\left(B^{d-1}\right)^{\perp} = Z_{d-1} = \ker\Delta^{-}$, and it is left to show that $\Delta^{-} f = |V| f$ for $f \in B^{d-1}$. Note that $B^{d-1} \subseteq Z^{d-1} = \ker\Delta_{X}^{+}$, and in addition, that since B^{d-1} only depends on X's $(d-1)$-skeleton,

$$B^{d-1}(X) = B^{d-1}\left(\overline{X}\right) \subseteq Z^{d-1}\left(\overline{X}\right) = \ker\Delta_{\overline{X}}^{+}.$$

Now from (1) it follows that for $f \in B^{d-1}$

$$\Delta_X^{-} f = \Delta_X^{-} f + \Delta_X^{+} f = \Delta_X f = |V| f - \Delta_{\overline{X}}^{+} f = |V| f$$

as desired. $\square$

The following proposition offers alternative characterizations of the spectral gap:

Proposition 3.3. *Let X be a d-dimensional complex with a complete skeleton.*

(1) The spectral gap of X is obtained by

$$\lambda(X) = \min\operatorname{Spec}\Delta. \tag{3.6}$$

(2) Furthermore, it is the $\binom{|V|-1}{d-1} + 1$ smallest eigenvalue of Δ^{+}.

Remarks.

(1) For graphs (3.6) gives $\lambda(G) = \min\operatorname{Spec}\left(\Delta^{+} + J\right)$, where $J = \Delta^{-} = \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & 1 & \cdots & 1 \\ \vdots & \vdots & \ddots & \vdots \\ 1 & 1 & \cdots & 1 \end{pmatrix}$.

(2) In general (3.6) does not hold: for example, for the triangle complex ⧓, $\lambda = \min\operatorname{Spec}\left(\Delta\Big|_{Z_{d-1}}\right) = 3$ but $\min\operatorname{Spec}\Delta = 1$.

Proof.

(1) First, since Δ decomposes w.r.t. $\Omega^{d-1} = B^{d-1} \oplus Z_{d-1}$ we have

$$\operatorname{Spec}\Delta = \operatorname{Spec}\Delta\Big|_{B^{d-1}} \bigcup \operatorname{Spec}\Delta\Big|_{Z_{d-1}} = \operatorname{Spec}\Delta^{-}\Big|_{B^{d-1}} \bigcup \operatorname{Spec}\Delta^{+}\Big|_{Z_{d-1}}.$$

By Proposition 3.2 $\operatorname{Spec}\Delta^{-}\Big|_{B^{d-1}} = \{|V|\}$ and $\operatorname{Spec}\Delta \subseteq [0, |V|]$, which implies that

$$\lambda = \min\operatorname{Spec}\left(\Delta^{+}\Big|_{Z_{d-1}}\right) = \min\operatorname{Spec}\Delta.$$

(2) The Euler characteristic satisfies $\sum_{i=-1}^{d} (-1)^{i} \left|X^{i}\right| = \chi(X) = \sum_{i=-1}^{d} (-1)^{i} \beta_i$. Therefore, by Proposition 2.2 we have $\lambda = \lambda_r$, with

$$\begin{aligned} r &= \left(\left|X^{d-1}\right| - \beta_{d-1}\right) - \left(\left|X^{d}\right| - \beta_{d}\right) \\ &= \left(\left|X^{d-1}\right| - \beta_{d-1}\right) - \left(\left|X^{d}\right| - \beta_{d}\right) + (-1)^{d} \sum_{i=-1}^{d} (-1)^{i} \left(\left|X^{i}\right| - \beta_{i}\right) \\ &= \sum_{i=-1}^{d-2} (-1)^{d+i} \left(\left|X^{i}\right| - \beta_{i}\right). \end{aligned}$$

Since the $(d-1)$-skeleton is complete, $\left|X^i\right| = \binom{|V|}{i+1}$ and $\beta_i = 0$ for $0 \le i \le d-2$, and so

$$r = \sum_{i=-1}^{d-2} (-1)^{d+i} \binom{|V|}{i+1} = \binom{|V|-1}{d-1}.$$

□

4 Proofs of the Main Theorems

4.1 The Cheeger Inequalities

This section is devoted to the proof of Theorem 1.2. The proof of the upper bound is a generalization of the standard proof for the Cheeger upper bound in graphs (1.3). The lower bound is comprised of three elements. The first is the Cheeger lower bound in graphs. The second is a connection between the Laplacian of a complex and the Laplacians of its one-dimensional links (see definition below). These connections were first shown by Garland [Gar73], and studied further by several authors [Zuk96, ABM05, GW12]. The third element is a connection between the combinatorial expansion of the complex and that of its links.

Proof of Theorem 1.2. We begin by proving the upper bound

$$\min \operatorname{Spec}\left(\Delta^+\Big|_{Z_{d-1}}\right) = \lambda(X) \le h(X) = \min_{V=\coprod_{i=0}^{d} A_i} \frac{|V| \cdot |F(A_0, A_1, \ldots, A_d)|}{|A_0| \cdot |A_1| \cdot \ldots \cdot |A_d|}.$$

Let $A_0, \ldots, A_d$ be a partition of V which realizes the minimum in h. We define $f \in \Omega^{d-1}$ by

$$f([\sigma_0 \, \sigma_1 \, \ldots \, \sigma_{d-1}]) = \begin{cases} \operatorname{sgn}(\pi) \left|A_{\pi(d)}\right| & \exists \pi \in \operatorname{Sym}_{\{0\ldots d\}} \text{ with } \sigma_i \in A_{\pi(i)} \text{ for } 0 \le i \le d-1 \\ 0 & \text{else, i.e. } \exists k, i \ne j \text{ with } \sigma_i, \sigma_j \in A_k. \end{cases} \tag{4.1}$$

Note that $f(\pi'\sigma) = \operatorname{sgn}(\pi') f(\sigma)$ for any $\pi' \in \operatorname{Sym}_{\{0\ldots d-1\}}$ and $\sigma \in X^{d-1}$. Therefore, f is a well-defined skew-symmetric function on oriented $(d-1)$-cells, i.e., $f \in \Omega^{d-1}$. Figure 3 illustrates f for $d = 1, 2$.



Figure 3: The form $f \in \Omega^{d-1}$ defined in (4.1), for complexes of dimensions one and two.

We proceed to show that $f \in Z_{d-1}$. Let $\sigma = [\sigma_0, \sigma_1, \ldots, \sigma_{d-2}] \in X_{\pm}^{d-2}$. As we assumed that X^{d-1} is complete,

$$(\partial_{d-1} f)(\sigma) = \sum_{v \sim \sigma} f([v, \sigma_0, \sigma_1, \ldots, \sigma_{d-2}]) = \sum_{v \notin \sigma} f([v, \sigma_0, \sigma_1, \ldots, \sigma_{d-2}]).$$

If for some k and $i \neq j$ we have $\sigma_i, \sigma_j \in A_k$, this sum vanishes. On the other hand, if there exists $\pi \in \operatorname{Sym}_{\{0\ldots d\}}$ such that $\sigma_i \in A_{\pi(i)}$ for $0 \leq i \leq d-2$ then

$$\begin{aligned}
(\partial_{d-1} f)(\sigma) &= \sum_{v \in A_{\pi(d-1)}} f([v, \sigma_0, \sigma_1, \ldots, \sigma_{d-2}]) + \sum_{v \in A_{\pi(d)}} f([v, \sigma_0, \sigma_1, \ldots, \sigma_{d-2}]) \\
&= \sum_{v \in A_{\pi(d-1)}} (-1)^{d-1} \operatorname{sgn} \pi \left|A_{\pi(d)}\right| + \sum_{v \in A_d} (-1)^d \operatorname{sgn} \pi \left|A_{\pi(d-1)}\right| \\
&= (-1)^{d-1} \operatorname{sgn} \pi \left(\left|A_{\pi(d-1)}\right| \left|A_{\pi(d)}\right| - \left|A_{\pi(d)}\right| \left|A_{\pi(d-1)}\right|\right) = 0
\end{aligned}$$

and in both cases $f \in Z_{d-1}$. Thus, by Rayleigh's principle

$$\lambda(X) = \min \operatorname{Spec}\left(\Delta\Big|_{Z_{d-1}}\right) \leq \frac{\langle \Delta f, f \rangle}{\langle f, f \rangle} = \frac{\langle \Delta^+ f, f \rangle}{\langle f, f \rangle} = \frac{\left\langle \partial_d^* f, \partial_d^* f \right\rangle}{\langle f, f \rangle}. \tag{4.2}$$

The denominator is

$$\langle f, f \rangle = \sum_{\sigma \in X^{d-1}} f(\sigma)^2,$$

and a $(d-1)$-cell σ contributes to this sum only if its vertices are in different blocks of the partition, i.e., there are no k and $i \neq j$ with $\sigma_i, \sigma_j \in A_k$. In this case, there exists a unique block, A_i, which does not contain a vertex of σ, and σ contributes $|A_i|^2$ to the sum. Since X^{d-1} is complete, there are $|A_0| \cdot \ldots \cdot |A_{i-1}| \cdot |A_{i+1}| \cdot \ldots \cdot |A_d|$ non-oriented $(d-1)$-cells whose vertices are in distinct blocks and which do not intersect A_i, hence

$$\langle f, f \rangle = \sum_{i=0}^{d} \left(\prod_{j \neq i} \left|A_j\right| \right) |A_i|^2 = |V| \prod_{i=0}^{d} |A_i|.$$

To evaluate the numerator in (4.2), we first show that for $\sigma \in X^d$

$$\left|(\partial_d^* f)(\sigma)\right| = \begin{cases} |V| & \sigma \in F(A_0, \ldots, A_d) \\ 0 & \sigma \notin F(A_0, \ldots, A_d) \end{cases}. \tag{4.3}$$

First, let $\sigma \notin F(A_0, \ldots, A_d)$. If σ has three vertices from the same A_i, or two pairs of vertices from the same blocks (i.e. $\sigma_i, \sigma_j \in A_k$ and $\sigma_{i'}, \sigma_{j'} \in A_{k'}$), then for every summand in

$$(\partial_d^* f)(\sigma) = \sum_{i=0}^{d} (-1)^i f(\sigma \backslash \sigma_i),$$

the cell $\sigma \backslash \sigma_i$ has two vertices from the same block, and therefore $\left(\partial_d^* f\right)(\sigma) = 0$. Next, assume that σ_j and σ_k (with $j < k$) is the only pair of vertices in σ which belong to the same block. The only non-vanishing terms in $\left(\partial_d^* f\right)(\sigma) = \sum_{i=0}^{d} (-1)^i f(\sigma \backslash \sigma_i)$ are $i = j$ and $i = k$, i.e.,

$$(\partial_d^* f)(\sigma) = (-1)^j f\left(\sigma \backslash \sigma_j\right) + (-1)^k f(\sigma \backslash \sigma_k).$$

Since the value of f on a simplex depends only on the blocks to which its vertices belong, we get

$$\begin{aligned}
f\left(\sigma \backslash \sigma_j\right) &= f\left(\left[\sigma_0\, \sigma_1\, \ldots\, \sigma_{j-1}\, \sigma_{j+1}\, \ldots \sigma_{k-1}\, \sigma_k\, \sigma_{k+1} \ldots\, \sigma_d\right]\right) \\
&= f\left(\left[\sigma_0\, \sigma_1\, \ldots\, \sigma_{j-1}\, \sigma_{j+1}\, \ldots \sigma_{k-1}\, \sigma_j\, \sigma_{k+1} \ldots\, \sigma_d\right]\right) \\
&= f\left((-1)^{k-j+1} \left[\sigma_0\, \sigma_1\, \ldots\, \sigma_{j-1}\, \sigma_j\, \sigma_{j+1}\, \ldots\, \sigma_{k-1}\, \sigma_{k+1} \ldots\, \sigma_d\right]\right) \\
&= (-1)^{k-j+1} f(\sigma \backslash \sigma_k),
\end{aligned}$$

so that

$$(\partial_d^* f)(\sigma) = (-1)^j (-1)^{k-j+1} f(\sigma\backslash\sigma_k) + (-1)^k f(\sigma\backslash\sigma_k) = 0.$$

The remaining case is $\sigma \in F(A_0, \ldots, A_d)$. Here, there exists $\pi \in \mathrm{Sym}_{\{0\ldots d\}}$ with $\sigma_i \in A_{\pi(i)}$ for $0 \le i \le d$. Observe that

$$f(\sigma\backslash\sigma_i) = \operatorname{sgn}(\pi \cdot (d\ d{-}1\ d{-}2\ \ldots\ i)) \left|A_{\pi(i)}\right| = (-1)^{d-i} \operatorname{sgn}\pi \left|A_{\pi(i)}\right|$$

and therefore

$$(\partial_d^* f)(\sigma) = \sum_{i=0}^{d} (-1)^i f(\sigma\backslash\sigma_i) = (-1)^d \operatorname{sgn}\pi \sum_{i=0}^{d} \left|A_{\pi(i)}\right| = (-1)^d \operatorname{sgn}\pi \left|V\right|,$$

and $\left|\left(\partial_d^* f\right)(\sigma)\right| = |V|$. This establishes (4.3), which implies that

$$\langle \partial_d^* f, \partial_d^* f \rangle = \sum_{\sigma \in X^d} \left|(\partial_d^* f)(\sigma)\right|^2 = |V|^2 \left|F(A_0, \ldots, A_d)\right|$$

and in total

$$\lambda(X) \le \frac{\left\langle \partial_d^* f, \partial_d^* f \right\rangle}{\langle f, f \rangle} = \frac{|V| \left|F(A_0, \ldots, A_d)\right|}{\prod_{i=0}^{d} |A_i|} = h(X),$$

settling the upper bound.

We proceed to prove the lower bound,

$$\frac{d\left(1 - \frac{d-1}{|V|}\right)^2}{8k} h^2(X) - (d-1)k \le \lambda(X).$$

For every $\tau \in X^{d-2}$ we consider the *link* of τ (see Figure 4),

$$\operatorname{lk}\tau = \left\{\sigma \in X \,\middle|\, \sigma \cap \tau = \varnothing \text{ and } \sigma \cup \tau \in X\right\}.$$



Figure 4: Two examples for the link of a vertex in a triangle complex.

Since $\dim\tau = d-2$, $\operatorname{lk}\tau$ is a graph, and there is a $1-1$ correspondence between vertices (edges) of $\operatorname{lk}\tau$ and $(d-1)$-cells (d-cells) of X which contain τ. As remarked above, we shall exploit connections between the Laplacian of X and those of $\operatorname{lk}\tau$. A summary of the properties which we shall use is given in Lemma A.1, in the appendix.

Assume $f \in Z_{d-1}$ is a normalized eigenfunction for $\lambda(X)$, i.e. $\langle f, f \rangle = 1$ and $\Delta^+ f = \lambda(X) f$. We observe the function f_τ on $(\operatorname{lk}\tau)^0$ (the vertices of $\operatorname{lk}\tau$) defined by

$$f_\tau(v) = f(v\tau).$$

The *local Laplacian of* X *at* τ, denoted by $\Delta_\tau^+ : \Omega^{d-1}(X) \to \Omega^{d-1}(X)$, is

$$\left(\Delta_\tau^+ f\right)(\sigma) = \begin{cases} \deg_\tau(\sigma) f(\sigma) - \sum\limits_{\substack{\sigma' \sim \sigma \\ \tau \subseteq \sigma'}} f(\sigma') & \tau \subset \sigma \\ 0 & \tau \nsubseteq \sigma \end{cases},$$

where $\deg_\tau(\sigma) = \#\left\{\sigma' \sim \sigma \,\middle|\, \tau \subseteq \sigma'\right\} = \deg_{\operatorname{lk}\tau}(\sigma \backslash \tau)$. Denoting by D the degree operator $(Df)(\sigma) = \deg(\sigma) f(\sigma)$, and letting $k = \max\left\{\deg \sigma \,\middle|\, \sigma \in X^{d-1}\right\} = \|D\|$, we obtain by Lemma A.1 (1), (2)

$$\begin{aligned} \lambda(X) = \langle \Delta^+ f, f \rangle &= \sum_{\tau \in X^{d-2}} \langle \Delta_\tau^+ f, f \rangle - (d-1) \langle Df, f \rangle \\ &= \sum_{\tau \in X^{d-2}} \langle \Delta_{\operatorname{lk}\tau}^+ f_\tau, f_\tau \rangle - (d-1) \langle Df, f \rangle \\ &\geq \sum_{\tau \in X^{d-1}} \langle \Delta_{\operatorname{lk}\tau}^+ f_\tau, f_\tau \rangle - (d-1) k. \end{aligned} \tag{4.4}$$

Point (3) in Lemma A.1 yields that $f_\tau \in Z_0(\operatorname{lk}\tau)$ for every $\tau \in X^{d-2}$. Denoting by $k_{\operatorname{lk}\tau}$ the maximal degree of a vertex in $\operatorname{lk}\tau$, the lower Cheeger bound for graphs (1.3) gives

$$\langle \Delta_{\operatorname{lk}\tau}^+ f_\tau, f_\tau \rangle \geq \lambda(\operatorname{lk}\tau) \langle f_\tau, f_\tau \rangle \geq \frac{h^2(\operatorname{lk}\tau)}{8 k_{\operatorname{lk}\tau}} \langle f_\tau, f_\tau \rangle \geq \frac{h^2(\operatorname{lk}\tau)}{8k} \langle f_\tau, f_\tau \rangle. \tag{4.5}$$

Combining (4.4) and (4.5) we find that

$$\lambda(X) \geq \sum_{\tau \in X^{d-2}} \frac{h^2(\operatorname{lk}\tau)}{8k} \langle f_\tau, f_\tau \rangle - (d-1) k. \tag{4.6}$$

In order to complete the proof, we use the following inequality which relates the local and global combinatorial expansions, and which will be proved afterwards:

$$h(\operatorname{lk}\tau) \geq \left(1 - \frac{d-1}{|V|}\right) h(X). \tag{4.7}$$

From (4.6) and (4.7) we have

$$\lambda(X) \geq \left(1 - \frac{d-1}{|V|}\right)^2 \frac{h^2(X)}{8k} \sum_{\tau \in X^{d-2}} \langle f_\tau, f_\tau \rangle - (d-1) k,$$

and therefore by point (4) in Lemma A.1

$$\lambda(X) \geq \left(1 - \frac{d-1}{|V|}\right)^2 \frac{d \cdot h^2(X)}{8k} - (d-1) k$$

which is the lower bound.

To prove (4.7), write $\tau = [\tau_0, \tau_1, \ldots, \tau_{d-2}]$ and denote $A_i = \{\tau_i\}$ for $0 \leq i \leq d-2$. Because of the correspondence between the edges and vertices of $\operatorname{lk}\tau$, and the d-cells and $(d-1)$-cells containing τ, we have

$$h(\operatorname{lk}\tau) \overset{def}{=} \min_{B \coprod C = (\operatorname{lk}\tau)^0} \frac{|E_{\operatorname{lk}\tau}(B, C)| \cdot \left|(\operatorname{lk}\tau)^0\right|}{|B| \cdot |C|} = \min_{B \coprod C = (\operatorname{lk}\tau)^0} \frac{|F(A_0, \ldots, A_{d-2}, B, C)| \cdot \left|(\operatorname{lk}\tau)^0\right|}{|B| \cdot |C|}.$$

Assume that the minimum is attained by $B = B_0$ and $C = C_0$. We define

$$A_{d-1} = B_0, \qquad A_d = V \backslash \left(\bigcup_{i=0}^{d-1} A_i \right).$$

Now $A_0, \ldots, A_d$ is a partition of V, and

$$F(A_0, \ldots, A_{d-2}, B_0, C_0) = F(A_0, \ldots, A_{d-2}, A_{d-1}, A_d)$$

since no d-cell containing τ has a vertex in $A_d \backslash C_0$. In addition,

$$\begin{aligned}
\frac{\left|(\operatorname{lk}\tau)^0\right| |A_d|}{|V| |C_0|} &\geq \frac{\left|(\operatorname{lk}\tau)^0\right| |A_d| - |A_{d-1}| (|A_d| - |C_0|)}{|V| |C_0|} \\
&= \frac{(|V| - (d-1) - (|A_d| - |C_0|)) |A_d| - |A_{d-1}| (|A_d| - |C_0|)}{|V| |C_0|} \\
&= \frac{(|V| - (d-1)) |A_d| - (|A_{d-1}| + |A_d|) (|A_d| - |C_0|)}{|V| |C_0|} \\
&= \frac{(|V| - (d-1)) [|A_d| - (|A_d| - |C_0|)]}{|V| |C_0|} = 1 - \frac{d-1}{|V|},
\end{aligned}$$

which implies

$$\begin{aligned}
h(\operatorname{lk}\tau) &= \frac{F(A_0, \ldots, A_{d-2}, A_{d-1}, A_d) \left|(\operatorname{lk}\tau)^0\right|}{|B_0| \cdot |C_0|} \\
&= \frac{F(A_0, \ldots, A_{d-2}, A_{d-1}, A_d) |V|}{|A_0| \cdot \ldots \cdot |A_d|} \cdot \frac{\left|(\operatorname{lk}\tau)^0\right| |A_d|}{|V| |C_0|} \\
&\geq h(X) \cdot \frac{\left|(\operatorname{lk}\tau)^0\right| |A_d|}{|V| |C_0|} \geq \left(1 - \frac{d-1}{|V|}\right) h(X).
\end{aligned}$$

□

4.2 The Mixing Lemma

Here we prove Theorem 1.4. We begin by formulating it precisely.

Theorem (1.4)**.** *Let X be a d-dimensional complex with a complete skeleton. Fix $\alpha \in \mathbb{R}$, and write* $\operatorname{Spec}(\alpha I - \Delta^+) = \{\rho_0 \geq \rho_1 \geq \ldots \geq \rho_m\}$*. For any disjoint sets of vertices $A_0, \ldots, A_d$ (not necessarily a partition), one has*

$$\left| |F(A_0, \ldots, A_d)| - \frac{\alpha \cdot |A_0| \cdot \ldots \cdot |A_d|}{|V|} \right| \leq \rho_\alpha \cdot (|A_0| \cdot \ldots \cdot |A_d|)^{\frac{d}{d+1}}$$

where

$$\rho_\alpha = \max \left\{ \left|\rho_{\binom{|V|-1}{d-1}}\right|, |\rho_m| \right\}.$$

Proof. For any disjoint sets of vertices $A_0, \ldots, A_{d-1}$, define $\delta_{A_0,\ldots,A_{d-1}} \in \Omega^{d-1}$ by

$$\delta_{A_0,\ldots,A_{d-1}}(\sigma) = \begin{cases} \operatorname{sgn}(\pi) & \exists \pi \in \operatorname{Sym}_{\{0\ldots d-1\}} \text{ with } \sigma_i \in A_{\pi(i)} \text{ for } 0 \le i \le d-1 \\ 0 & \text{else} \end{cases}.$$

Since the skeleton of X is complete,

$$\|\delta_{A_0,\ldots,A_{d-1}}\| = \sqrt{\sum_{\sigma \in X^{d-1}} \delta^2_{A_0,\ldots,A_{d-1}}(\sigma)} = \sqrt{|A_0| \cdot \ldots \cdot |A_{d-1}|}. \tag{4.8}$$

Now, let $A_0, \ldots, A_d$ be disjoint subsets of V (not necessarily a partition), and denote

$$\varphi = \delta_{A_0,A_1,A_2,\ldots,A_{d-1}}$$
$$\psi = \delta_{A_d,A_1,A_2,\ldots,A_{d-1}}.$$

Let σ be an oriented $(d-1)$-cell with one vertex in each of $A_0, A_1, \ldots, A_{d-1}$. With some abuse of notation we shall denote this by $\sigma \in F(A_0, \ldots, A_{d-1})$, ignoring the orientation of σ. There is a correspondence between d-cells in $F(A_0, \ldots, A_d)$ containing σ, and neighbors of σ which lie in $F(A_d, A_1, \ldots, A_{d-1})$. Furthermore, for such a neighbor σ' we have $\varphi(\sigma) = \psi(\sigma')$, since σ and σ' must share the vertices which belong to $A_1, \ldots, A_{d-1}$. Therefore, by (3.2)

$$\begin{aligned} \langle \varphi, (D - \Delta^+)\psi \rangle &= \sum_{\sigma \in X^{d-1}} \varphi(\sigma)\left((D - \Delta^+)\psi\right)(\sigma) = \sum_{\sigma \in X^{d-1}} \sum_{\sigma' \sim \sigma} \varphi(\sigma)\psi(\sigma') \\ &= \sum_{\sigma \in F(A_0 \ldots A_{d-1})} \sum_{\sigma' \sim \sigma} \varphi(\sigma)\psi(\sigma') = \sum_{\sigma \in F(A_0 \ldots A_{d-1})} \#\left\{\sigma' \in F(A_d, A_1, \ldots, A_{d-1}) \,\middle|\, \sigma' \sim \sigma\right\} \\ &= \sum_{\sigma \in F(A_0 \ldots A_{d-1})} \#\left\{\tau \in F(A_0, A_1, \ldots, A_d) \,\middle|\, \sigma \subseteq \tau\right\} = |F(A_0, A_1, \ldots, A_d)|. \end{aligned} \tag{4.9}$$

Notice that since the A_i are disjoint, φ and ψ are supported on different $(d-1)$-cells, so that for any $\alpha \in \mathbb{R}$

$$\langle \varphi, (D - \Delta^+)\psi \rangle = \langle \varphi, -\Delta^+\psi \rangle = \langle \varphi, (\alpha I - \Delta^+)\psi \rangle. \tag{4.10}$$

By the orthogonal decomposition $\Omega^{d-1} = B^{d-1} \oplus Z_{d-1}$ and the fact that $B^{d-1} \subseteq Z^{d-1} = \ker \Delta^+$,

$$\begin{aligned} |F(A_0, A_1, \ldots, A_d)| &= \langle \varphi, (\alpha I - \Delta^+)\psi \rangle \\ &= \langle \varphi, (\alpha I - \Delta^+)(\mathbb{P}_{B^{d-1}}\psi + \mathbb{P}_{Z_{d-1}}\psi) \rangle \\ &= \langle \varphi, \alpha \mathbb{P}_{B^{d-1}}\psi + (\alpha I - \Delta^+)\mathbb{P}_{Z_{d-1}}\psi \rangle \\ &= \alpha \langle \varphi, \mathbb{P}_{B^{d-1}}\psi \rangle + \langle \varphi, (\alpha I - \Delta^+)\mathbb{P}_{Z_{d-1}}\psi \rangle. \end{aligned} \tag{4.11}$$

We proceed to evaluate each of these terms separately. Using (3.5) and (3.4) we find that

$$\alpha \langle \varphi, \mathbb{P}_{B^{d-1}}\psi \rangle = \frac{\alpha}{|V|} \langle \varphi, \Delta^-\psi \rangle = \frac{\alpha}{|V|} \left\langle \varphi, \left(|V| I - \Delta^+_X - \Delta^+_{\overline{X}}\right)\psi \right\rangle$$

and by (4.9) and (4.10) this implies

$$\begin{aligned} \alpha \langle \varphi, \mathbb{P}_{B^{d-1}}\psi \rangle &= \frac{\alpha}{|V|} \langle \varphi, |V| I - \Delta^+_X \psi \rangle + \frac{\alpha}{|V|} \left\langle \varphi, -\Delta^+_{\overline{X}}\psi \right\rangle \\ &= \frac{\alpha}{|V|} |F_X(A_0, A_1, \ldots, A_d)| + \frac{\alpha}{|V|} \left|F_{\overline{X}}(A_0, A_1, \ldots, A_d)\right| \\ &= \frac{\alpha \cdot |A_0| \cdot \ldots \cdot |A_d|}{|V|}. \end{aligned} \tag{4.12}$$

We turn to the second term in (4.11). First, we recall from the proof of Proposition 3.3 that $\dim B^{d-1} = \binom{|V|-1}{d-1}$. Since $B^{d-1} \subseteq \ker \Delta^{+}$, we have that in $\operatorname{Spec}\left(\alpha I - \Delta^{+}\right) = \left\{\rho_0 \geq \rho_1 \geq \ldots \geq \rho_m\right\}$ the first $\binom{|V|-1}{d-1}$ values correspond to B^{d-1}, and the rest to $\left(B^{d-1}\right)^{\perp} = Z_{d-1}$. Thus,

$$\rho_\alpha = \max\left\{\left|\rho_{\binom{|V|-1}{d-1}}\right|, \left|\rho_m\right|\right\} = \max\left\{|\lambda| \,\middle|\, \lambda \in \operatorname{Spec}\left(\alpha I - \Delta\right)\big|_{Z_{d-1}}\right\} = \left\|\left(\alpha I - \Delta\right)\big|_{Z_{d-1}}\right\|, \tag{4.13}$$

and therefore

$$\begin{aligned} \left|\left\langle \varphi, \left(\alpha I - \Delta^{+}\right)\mathbb{P}_{Z_{d-1}}\psi\right\rangle\right| &\leq \|\varphi\| \cdot \left\|\left(\alpha I - \Delta^{+}\right)\mathbb{P}_{Z_{d-1}}\psi\right\| \leq \|\varphi\| \cdot \left\|\left(\alpha I - \Delta^{+}\right)\big|_{Z_{d-1}}\right\| \cdot \left\|\mathbb{P}_{Z_{d-1}}\psi\right\| \\ &\leq \rho_\alpha \cdot \|\varphi\| \cdot \|\psi\| = \rho_\alpha \sqrt{|A_0|\,|A_d|}\,|A_1|\,|A_2|\ldots|A_{d-1}|, \end{aligned} \tag{4.14}$$

where the last step is by (4.8). Together (4.11), (4.12) and (4.14) give

$$\left|\left|F\left(A_0, A_1, \ldots, A_d\right)\right| - \frac{\alpha \cdot |A_0| \cdot \ldots \cdot |A_d|}{|V|}\right| \leq \rho_\alpha \sqrt{|A_0|\,|A_d|}\,|A_1|\,|A_2|\ldots|A_{d-1}|.$$

Since $A_0, \ldots, A_d$ play the same role, one can also obtain the bound

$$\rho_\alpha \sqrt{\left|A_{\pi(0)}\right|\left|A_{\pi(d)}\right|}\left|A_{\pi(1)}\right|\left|A_{\pi(2)}\right|\ldots\left|A_{\pi(d-1)}\right|,$$

for any $\pi \in \operatorname{Sym}_{\{0..d\}}$. Taking the geometric mean over all such π gives

$$\left|\left|F\left(A_0, A_1, \ldots, A_d\right)\right| - \frac{\alpha \cdot |A_0| \cdot \ldots \cdot |A_d|}{|V|}\right| \leq \rho_\alpha \cdot \left(|A_0|\,|A_1|\ldots|A_d|\right)^{\frac{d}{d+1}}.$$

□

Remark. The estimate (4.14) is somewhat wasteful. As is done in graphs, a slightly better one is

$$\left|\left\langle \varphi, \left(\alpha I - \Delta^{+}\right)\mathbb{P}_{Z_{d-1}}\psi\right\rangle\right| = \left|\left\langle \mathbb{P}_{Z_{d-1}}\varphi, \left(\alpha I - \Delta^{+}\right)\mathbb{P}_{Z_{d-1}}\psi\right\rangle\right| \leq \rho_\alpha \cdot \left\|\mathbb{P}_{Z_{d-1}}\varphi\right\| \cdot \left\|\mathbb{P}_{Z_{d-1}}\psi\right\|,$$

and we leave it to the curious reader to verify that this gives

$$\left|\left\langle \varphi, \left(\alpha I - \Delta^{+}\right)\mathbb{P}_{Z_{d-1}}\psi\right\rangle\right| \leq \rho_\alpha \sqrt{|A_0|\left(1 - \frac{\sum_{i=0}^{d-1}|A_i|}{|V|}\right)|A_d|\left(1 - \frac{\sum_{i=1}^{d}|A_i|}{|V|}\right)}\,|A_1|\ldots|A_{d-1}|.$$

4.3 Gromov's Geometric Overlap

Here we prove Corollary 1.6, which gives a bound on the geometric overlap of a complex in terms of the width of its spectrum.

Proof of Corollary 1.6. Given $\varphi : V \to \mathbb{R}^{d+1}$, choose arbitrarily some partition of V into equally sized parts $P_0, \ldots, P_d$. By Pach's theorem [Pac98], there exist $c_d > 0$ and $Q_i \subseteq P_i$ of sizes $|Q_i| \geq c_d |P_i|$ such that for some $x \in \mathbb{R}^{d+1}$ we have $x \in \operatorname{conv}\left\{\varphi(v) \,\middle|\, v \in \sigma\right\}$ for any $\sigma \in F\left(Q_0, \ldots, Q_d\right)$. Denote

$$\lambda_{\min} = \lambda(X) = \min \operatorname{Spec} \Delta\Big|_{Z_{d-1}}, \quad \lambda_{\max} = \max \operatorname{Spec} \Delta\Big|_{Z_{d-1}}.$$

By the Mixing Lemma (Theorem 1.4), with $\alpha = \frac{1}{2}(\lambda_{\max} + \lambda_{\min})$, we have

$$\begin{aligned} |F(Q_0, \ldots, Q_d)| &\geq \frac{\alpha \cdot |Q_0| \cdot \ldots \cdot |Q_d|}{|V|} - \rho_\alpha \cdot (|Q_0| \cdot \ldots \cdot |Q_d|)^{\frac{d}{d+1}} \\ &\geq \left(\frac{c_d |V|}{d+1}\right)^d \left(\frac{\alpha c_d}{d+1} - \rho_\alpha\right) \\ &= \left(\frac{c_d |V|}{d+1}\right)^d \left(\frac{(\lambda_{\max} + \lambda_{\min}) c_d}{2d+2} - \frac{\lambda_{\max} - \lambda_{\min}}{2}\right) \\ &= \left(\frac{c_d |V|}{d+1}\right)^d \left(\frac{1}{2} + \frac{c_d}{2d+2}\right) \left(\frac{\lambda_{\min}}{\lambda_{\max}} - \frac{d+1-c_d}{d+1+c_d}\right) \lambda_{\max}. \end{aligned}$$

Choosing $\varepsilon_d = 1 - \frac{d+1-c_d}{d+1+c_d}$, and recalling that

$$1 - \varepsilon_d < 1 - \varepsilon < \frac{\lambda_{\min}}{\lambda_{\max}}$$

we get

$$|F(Q_0, \ldots, Q_d)| \geq \left(\frac{c_d |V|}{d+1}\right)^d \left(\frac{1}{2} + \frac{c_d}{2d+2}\right) (\varepsilon_d - \varepsilon) \lambda_{\max}.$$

Since this holds for any $\varphi : V \to \mathbb{R}^{d+1}$,

$$\operatorname{overlap}(X) \geq \underbrace{\left(\frac{c_d}{d+1}\right)^d \left(\frac{1}{2} + \frac{c_d}{2d+2}\right) (\varepsilon_d - \varepsilon)}_{\delta} \frac{|V|^d}{|X^d|} \lambda_{\max}.$$

□

4.4 Expansion in random complexes

In this section we prove Corollaries 1.3 and 1.7. The main idea is the following Lemma, which is a variation on the analysis in [GW12] of the spectrum of $D - \Delta^+$ for $X = X(d, n, p)$.

Lemma 4.1. *Let $c > 0$. For $C > C_0 = C_0(c, d)$ there exists $\gamma = \gamma(C)$ satisfying $0 < \gamma < C$ and $\gamma = o(C)$ such that $X = X\left(d, n, \frac{C \cdot \log n}{n}\right)$ satisfies*

$$\operatorname{Spec}\left(\Delta^+\Big|_{Z_{d-1}}\right) \subseteq [(C-\gamma)\log n, (C+\gamma)\log n]$$

with probability at least $1 - n^{-c}$.

Proof. We denote $p = \frac{C \cdot \log n}{n}$. For C large enough we shall find γ with the desired properties such that

$$\left\| \left(\Delta_X^+ - pn \cdot I\right)\Big|_{Z_{d-1}} \right\| \leq \gamma \log n \tag{4.15}$$

holds with probability at least $1 - n^{-c}$. Whenever (4.15) holds Weyl's perturbation inequality[5] gives

$$\operatorname{Spec}\left(\Delta^+\Big|_{Z_{d-1}}\right) \subseteq [pn - \gamma \log n, pn + \gamma \log n] = [(C-\gamma)\log n, (C+\gamma)\log n],$$

---

[5] In fact, we need only a corollary of it, which is that $\max \operatorname{Spec}(A+B) \leq \max(\operatorname{Spec} A) + \|B\|$ for any hermitian operators A, B, and the same for $\min \operatorname{Spec}$.

and the Lemma follows.

To show (4.15) we use

$$\begin{aligned}\left\|\left(\Delta_X^+ - pn\cdot I\right)\Big|_{Z_{d-1}}\right\| &= \left\|\left(\Delta_X^+ - (n-d)\,pI - pdI + D - D\right)\Big|_{Z_{d-1}}\right\| \\ &\le \left\|\left(D - (n-d)\,pI\right)\Big|_{Z_{d-1}}\right\| + \left\|\left(D - \Delta_X^+ + pdI\right)\Big|_{Z_{d-1}}\right\| \qquad (4.16)\end{aligned}$$

and we will treat each term separately. For the first, we have

$$\left\|\left(D - (n-d)\,pI\right)\Big|_{Z_{d-1}}\right\| \le \left\|D - (n-d)\,pI\right\| = \max_{\sigma\in X^{d-1}} \left|\deg\sigma - (n-d)\,p\right|.$$

Since $\deg\sigma \sim B\,(n-d,p)$, a Chernoff type bound (e.g. [Jan02, Theorem 1]) gives that for every $t>0$

$$\mathrm{Prob}\left(\left|\deg\sigma - (n-d)\,p\right| > t\right) \le 2e^{-\frac{t^2}{2(n-d)p+\frac{2t}{3}}}.$$

By a union bound on the degrees of the $(d-1)$-cells we get

$$\mathrm{Prob}\left(\max_{\sigma\in X^{d-1}} \left|\deg\sigma - (n-d)\,p\right| > t\right) \le 2\binom{n}{d} e^{-\frac{t^2}{2(n-d)p+\frac{2t}{3}}}, \qquad (4.17)$$

and a straightforward calculation shows that there exists $\alpha = \alpha\,(c,d) > 0$ such that for $t = \alpha\sqrt{np\log n}$, the r.h.s. in (4.17) is bounded by $\frac{1}{2n^c}$ for large enough C and n. In total this implies

$$\mathrm{Prob}\left(\left\|\left(D - (n-d)\,pI\right)\Big|_{Z_{d-1}}\right\| \le \alpha\sqrt{C}\log n\right) \ge 1 - \frac{1}{2n^c}.$$

Turning to the last term in (4.16), we go back to the estimates in the proof of Theorem 7 in [GW12] and in Corollary 7.1 in [Oli10]. Denoting by D_{comp} and Δ^+_{comp} the degree operator and upper Laplacian of the full d-complex on n vertices, we have $\left(D_{comp} - \Delta^+_{comp}\right)\Big|_{Z_{d-1}} \equiv -d\cdot I$, and thus by these estimates

$$\mathrm{Prob}\left(\left\|\left(D - \Delta_X^+\right)\Big|_{Z_{d-1}} + pdI\Big|_{Z_{d-1}}\right\| \ge t\right) \le 2\binom{n}{d} e^{-\frac{t^2}{8pnd+4t}}.$$

Again, there exists $\beta = \beta\,(c,d) > 0$ such that for $t = \beta\sqrt{np\log n}$, the r.h.s. is bounded by $\frac{1}{2n^c}$ for large enough C and n. Consequently,

$$\mathrm{Prob}\left(\left\|\left(D - \Delta_X^+\right)\Big|_{Z_{d-1}}\right\| \le \beta\sqrt{C}\log n\right) \ge 1 - \frac{1}{2n^c},$$

so that

$$\mathrm{Prob}\left(\left\|\left(\Delta_X^+ - pnI\right)\Big|_{Z_{d-1}}\right\| \le (\alpha+\beta)\,\sqrt{C}\log n\right) \ge 1 - n^{-c},$$

and $\gamma = (\alpha+\beta)\,\sqrt{C}$ gives the required result. □

We obtain the following corollary, which is the content of Corollaries 1.3 and 1.7.

Corollary 4.2. *Fix* $c > 0$*. There exist a constant* $H = H(C) > 0$ *satisfying* $\lim_{C\to\infty} \frac{H}{C} = 1$*, and a constant* $\mathcal{O} = \mathcal{O}_d > 0$*, such that for* $C \geq C_0(c,d)$ *and* $n \geq n_0(c,d)$

$$\operatorname{Prob}\left(h\left(X\left(d,n,\frac{C\cdot\log n}{n}\right)\right) \geq H\cdot\log n\right) \geq 1 - n^{-c} \tag{4.18}$$

and

$$\operatorname{Prob}\left(\operatorname{overlap}\left(X\left(d,n,\frac{C\cdot\log n}{n}\right)\right) > \mathcal{O}\right) \geq 1 - n^{-c}.$$

Proof. Since $\lambda(X) \leq h(X)$ (Theorem 1.2), (4.18) follows from Lemma 4.1 with $H = C - \gamma$. We turn to the geometric overlap. Fix some ε smaller than ε_d from Corollary 1.6. From Lemma 4.1 it follows that for C large enough

$$\frac{\lambda_{\min}(X)}{\lambda_{\max}(X)} \geq \frac{C-\gamma}{C+\gamma} > 1 - \varepsilon \tag{4.19}$$

with probability at least $1 - n^{-c}$. When this holds, by Corollary 1.6 there exists $\delta > 0$ such that

$$\operatorname{overlap}(X) > \delta\cdot\frac{|V|^d}{|X^d|}\cdot\lambda_{\max}(X) \geq \delta\cdot\frac{n^d}{|X^d|}\cdot(C-\gamma)\log n.$$

Since $\left|X^d\right| \sim B\left(\binom{n}{d+1},\frac{C\log n}{n}\right)$ by Chernoff's bound there exists $C' > 0$ such that

$$\left|X^d\right| \leq \frac{C\log n}{n}\binom{n}{d+1} + C'\log n\sqrt{\frac{C}{n}\binom{n}{d+1}} \tag{4.20}$$

with probability at least $1 - n^{-c}$. When both (4.19) and (4.20) hold,

$$\operatorname{overlap}(X) > \delta\cdot\frac{n^d}{|X^d|}\cdot(C-\gamma)\log n \geq \frac{\delta n^d (C-\gamma)}{\frac{C}{n}\binom{n}{d+1} + C'\sqrt{\frac{C}{n}\binom{n}{d+1}}} \xrightarrow{n\to\infty} \delta(d+1)!\left(1 - \frac{\gamma}{C}\right),$$

so that the conclusion holds for any $\mathcal{O} < \delta(d+1)!$. □

5 Open Questions

Cheeger inequality and mixing without a complete skeleton

The proofs for the upper bound in the generalized Cheeger inequality, and that of the generalized mixing lemma, assume that the skeleton is complete. This of course raises the following questions:

Question: Can one find a good upper bound for $\lambda(X)$ by $h(X)$ for general simplicial complexes?

Question: Can the discrepancy in X be bounded for general simplicial complexes?

Tightness of the bounds in the generalized Cheeger inequalities

For the complete d-dimensional simplicial complex, $h(X) = |V| = \lambda(X)$. This shows that the upper bound on $\lambda(X) \leq h(X)$ is tight.

Question: Is the lower bound $\frac{d\left(1-\frac{d-1}{|V|}\right)^2}{8k}h^2(X) - (d-1)k \leq \lambda(X)$ asymptotically tight for $d \geq 2$?

Inverse Mixing Lemma

In [BL06] Bilu and Linial proved an Inverse Mixing Lemma for graphs:

Theorem ([BL06])**.** *Let G be a d-regular graph on n vertices. Suppose that for any $A, B \subseteq V$, with $A \cap B = \varnothing$*

$$\left| E(A,B) - \frac{|A|\,|B|\,d}{n} \right| \leq \alpha \sqrt{|A|\,|B|}.$$

Then all but the largest eigenvalue of G are bounded, in absolute value, by $O\left(\alpha\left(1 + \log\left(\frac{d}{\alpha}\right)\right)\right)$.

Question: Can one prove a generalized Inverse Mixing Lemma for simplicial complexes?

Random simplicial complexes

In the random graph model $G = G(n,p) = X(1,n,p)$, taking $p = \frac{k}{n}$ with a fixed k gives disconnected G a.a.s. However, random k-regular graphs are a.a.s. connected, and in fact are excellent expanders (see e.g. [HLW06]). In dimension d, the $(d-1)$-homology of $X(d,n,p)$ is nontrivial a.a.s. for $p = \frac{k}{n}$, and it is natural to ask about the expansion quality of regular complexes. Since it is not clear whether such complexes even exist, we say that a *k-semiregular* complex is a complex with $k - \sqrt{k} \leq \deg \sigma \leq k + \sqrt{k}$ for all $\sigma \in X^{\dim X - 1}$, and ask:

Question: Are $h(X)$ and $\operatorname{overlap}(X)$ bounded away from zero for X a random k-semiregular d-complex?

A Riemannian Analogue

In Riemannian geometry, the Cheeger constant of a Riemannian manifold M is concerned with its partitions into two submanifolds along a common boundary of codimension one. The original Cheeger inequalities, due to Cheeger [Che70] and Buser [Bus82], relate the Cheeger constant to the smallest eigenvalue of the Laplace-Beltrami on $C^{\infty}(M) = \Omega^0(M)$.

Question: Can one define an isoperimetric quantity which concerns partitioning of M into $d+1$ parts, and relate it to the spectrum of the Laplace-Beltrami operator on $\Omega^{d-1}(M)$, the space of differential $(d-1)$-forms?

Ramanujan Complexes

Ramanujan Graphs are expanders which are spectrally optimal in the sense of the Alon-Boppana theorem [Nil91], and therefore excellent combinatorial expanders. Such graphs were constructed in [LPS88] as quotients of the Bruhat-Tits tree associated with $\mathrm{PSL}_2\left(\mathbb{Q}_p\right)$ by certain arithmetic lattices. Analogue quotients of the Bruhat-Tits buildings associated with $\mathrm{PSL}_n\left(\mathbb{F}_q(t)\right)$ are constructed in [LSV05], and termed *Ramanujan Complexes*. It is natural to ask whether these complexes are also optimal expanders in the spectral and combinatorial senses.

A Appendix

The following lemma, which already appeared in [GW12] (for a normalized version of the Laplacian), lists some connections between the upper Laplacian of a complex and those of its links. For completeness we give a proof here.

Lemma A.1. *Let X be a d-dimensional simplicial complex, and for $f \in \Omega^{d-1}, \sigma \in X^{d-1}, \tau \in X^{d-2}$ define*

$$\begin{aligned}
(Df)(\sigma) &= \deg(\sigma) f(\sigma) \\
\deg_\tau(\sigma) &= \deg_\tau(\sigma) = \#\left\{\sigma' \sim \sigma \,\middle|\, \tau \subseteq \sigma'\right\} = \deg_{\operatorname{lk}\tau}(\sigma \backslash \tau) \\
(\Delta_\tau^+ f)(\sigma) &= \begin{cases} \deg_\tau(\sigma) f(\sigma) - \sum\limits_{\substack{\sigma' \sim \sigma \\ \tau \subseteq \sigma'}} f(\sigma') & \tau \subset \sigma \\ 0 & \tau \nsubseteq \sigma \end{cases} \\
f_\tau(v) &= f(v\tau).
\end{aligned}$$

The following then hold:

(1) $\Delta^+ = \sum\limits_{\tau \in X^{d-2}} \Delta_\tau^+ - (d-1) D$.

(2) $\langle \Delta_\tau^+ f, f \rangle = \left\langle \Delta_{\operatorname{lk}\tau}^+ f_\tau, f_\tau \right\rangle$.

(3) If $f \in Z_{d-1}$ *then* $f_\tau \in Z_0(\operatorname{lk}\tau)$.

(4) $\sum\limits_{\tau \in X^{d-2}} \langle f_\tau, f_\tau \rangle = d \langle f, f \rangle$.

Proof.

(1) By the definition of Δ_τ^+,

$$\begin{aligned}
\sum_{\tau \in X^{d-2}} \Delta_\tau^+ f(\sigma) - (d-1) Df(\sigma) &= \sum_{\substack{\tau \in X^{d-2} \\ \tau \subseteq \sigma}} \left(\deg_\tau(\sigma) f(\sigma) - \sum_{\substack{\sigma' \sim \sigma \\ \tau \subseteq \sigma'}} f\left(\sigma'\right) \right) - (d-1) \deg(\sigma) f(\sigma) \\
&= \left(\sum_{\substack{\tau \in X^{d-2} \\ \tau \subseteq \sigma}} \deg_\tau(\sigma) - (d-1) \deg \sigma f(\sigma) \right) - \sum_{\substack{\tau \in X^{d-2} \\ \tau \subseteq \sigma}} \sum_{\substack{\sigma' \sim \sigma \\ \tau \subseteq \sigma'}} f\left(\sigma'\right) \\
&= \deg(\sigma) f(\sigma) - \sum_{\sigma' \sim \sigma} f\left(\sigma'\right) = \Delta^+ f(\sigma).
\end{aligned}$$

(2) Let $f \in \Omega^{d-1}$ and $\tau \in X^{d-2}$. We first notice that $(\Delta_\tau^+ f)_\tau = \Delta_{\operatorname{lk}\tau}^+ f_\tau$, since

$$\begin{aligned}
(\Delta_\tau^+ f)_\tau(v) &= (\Delta_\tau^+ f)(v\tau) = \deg_\tau(v\tau) f(v\tau) - \sum_{\substack{\sigma' \sim v\tau \\ \tau \subseteq \sigma'}} f\left(\sigma'\right) \\
&= \deg_\tau(v\tau) f(v\tau) - \sum_{\substack{v' \sim \tau \\ v'\tau \sim v\tau}} f\left(v'\tau\right) = \deg_{\operatorname{lk}\tau}(v) f_\tau(v) - \sum_{v' \underset{\operatorname{lk}\tau}{\sim} v} f_\tau\left(v'\right) = \Delta_{\operatorname{lk}\tau}^+ f_\tau(v).
\end{aligned}$$

Since $(\operatorname{lk}\tau)^0 = \{v \in V \,|\, v \sim \tau\}$, this gives

$$\langle \Delta_{\operatorname{lk}\tau}^+ f_\tau, f_\tau \rangle = \left\langle (\Delta_\tau^+ f)_\tau, f_\tau \right\rangle = \sum_{v \sim \tau} (\Delta_\tau^+ f)_\tau(v) f_\tau(v) = \sum_{v \sim \tau} (\Delta_\tau^+ f)(v\tau) f(v\tau) = \langle \Delta_\tau^+ f, f \rangle$$

where the last equality is since $\Delta_\tau^+ f$ is supported on $(d-1)$-cells containing τ.

(3) If $f \in Z_{d-1}$ and $\tau \in X^{d-2}$ then

$$\left(\partial_0^{\operatorname{lk}\tau} f_\tau\right)(\varnothing) = \sum_{v \in (\operatorname{lk}\tau)^0} f_\tau(v) = \sum_{v \in (\operatorname{lk}\tau)^0} f(v\tau) = \sum_{v \sim \tau} f(v\tau) = (\partial_{d-1} f)(\tau) = 0$$

implies that $f_\tau \in Z_0(\operatorname{lk}\tau)$.

(4) This is by

$$\sum_{\tau \in X^{d-2}} \langle f_\tau, f_\tau \rangle = \sum_{\tau \in X^{d-2}} \sum_{v \sim \tau} f_\tau^2(v) = \sum_{\tau \in X^{d-2}} \sum_{v \sim \tau} f^2(v\tau) = d \sum_{\sigma \in X^{d-1}} f^2(\sigma) = d \langle f, f \rangle .$$

□

References

[ABM05] R. Aharoni, E. Berger, and R. Meshulam. Eigenvalues and homology of flag complexes and vector representations of graphs. *Geometric and functional analysis*, 15(3):555–566, 2005.

[AC88] N. Alon and F.R.K. Chung. Explicit construction of linear sized tolerant networks. *Discrete Mathematics*, 72(1-3):15–19, 1988.

[Alo86] N. Alon. Eigenvalues and expanders. *Combinatorica*, 6(2):83–96, 1986.

[AM85] N. Alon and V.D. Milman. λ_1, isoperimetric inequalities for graphs, and superconcentrators. *Journal of Combinatorial Theory, Series B*, 38(1):73–88, 1985.

[BL06] Y. Bilu and N. Linial. Lifts, discrepancy and nearly optimal spectral gap. *Combinatorica*, 26(5):495–519, 2006.

[Bus82] P. Buser. A note on the isoperimetric constant. *Ann. Sci. École Norm. Sup.(4)*, 15(2):213–230, 1982.

[Che70] J. Cheeger. A lower bound for the smallest eigenvalue of the laplacian. *Problems in analysis*, 195:199, 1970.

[Chu97] F.R.K. Chung. *Spectral graph theory*. Number 92. Amer Mathematical Society, 1997.

[Dod84] J. Dodziuk. Difference equations, isoperimetric inequality and transience of certain random walks. *Trans. Amer. Math. Soc*, 284, 1984.

[Eck44] B. Eckmann. Harmonische funktionen und randwertaufgaben in einem komplex. *Commentarii Mathematici Helvetici*, 17(1):240–255, 1944.

[FGL$^+$11] J. Fox, M. Gromov, V. Lafforgue, A. Naor, and J. Pach. Overlap properties of geometric expanders. In *Proceedings of the Twenty-Second Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1188–1197. SIAM, 2011.

[Gar73] H. Garland. p-adic curvature and the cohomology of discrete subgroups of p-adic groups. *The Annals of Mathematics*, 97(3):375–423, 1973.

[Gro10] M. Gromov. Singularities, expanders and topology of maps. part 2: From combinatorics to topology via algebraic isoperimetry. *Geometric And Functional Analysis*, 20(2):416–526, 2010.

[GW12] A. Gundert and U. Wagner. On laplacians of random complexes. 2012.

[HLW06] S. Hoory, N. Linial, and A. Wigderson. Expander graphs and their applications. *Bulletin of the American Mathematical Society*, 43(4):439–562, 2006.

[Jan02] S. Janson. On concentration of probability. *Contemporary combinatorics*, 10(3):1–9, 2002.

[LM06] N. Linial and R. Meshulam. Homological connectivity of random 2-complexes. *Combinatorica*, 26(4):475–487, 2006.

[LPS88] A. Lubotzky, R. Phillips, and P. Sarnak. Ramanujan graphs. *Combinatorica*, 8(3):261–277, 1988.

[LSV05] A. Lubotzky, B. Samuels, and U. Vishne. Ramanujan complexes of type $\tilde{A}_d$. *Israel Journal of Mathematics*, 149(1):267–299, 2005.

[Lub12] A. Lubotzky. Expander graphs in pure and applied mathematics. *Bull. Amer. Math. Soc*, 49:113–162, 2012.

[MW11] J. Matoušek and U. Wagner. On gromov's method of selecting heavily covered points. *Arxiv preprint arXiv:1102.3515*, 2011.

[Nil91] A. Nilli. On the second eigenvalue of a graph. *Discrete Mathematics*, 91(2):207–210, 1991.

[NR12] Ilan Newman and Yuri Rabinovich. On multiplicative λ-approximations and some geometric applications. In *Proceedings of the Twenty-Third Annual ACM-SIAM Symposium on Discrete Algorithms*, SODA '12, pages 51–67. SIAM, 2012.

[Oli10] R.I. Oliveira. Concentration of the adjacency matrix and of the laplacian in random graphs with independent edges. *Arxiv preprint ArXiv:0911.0600v2*, 2010.

[Pac98] J. Pach. A tverberg-type result on multicolored simplices. *Computational Geometry*, 10(2):71–76, 1998.

[PR12] O. Parzanchevski and R. Rosenthal. Random walks on simplicial complexes. *Preprint*, 2012.

[Tan84] R.M. Tanner. Explicit concentrators from generalized n-gons. *SIAM Journal on Algebraic and Discrete Methods*, 5:287, 1984.

[Tao11] T. Tao. Basic theory of expander graphs. `http://terrytao.wordpress.com/2011/12/02/245b-notes-1-basic-theory-of-expander-graphs/`, 2011.

[Zuk96] A. Zuk. La propriété (T) de Kazhdan pour les groupes agissant sur les polyedres. *Comptes rendus de l'Académie des sciences. Série 1, Mathématique*, 323(5):453–458, 1996.